

## Depression in Women

Contrary to popular belief, clinical depression is not a “normal part of being a woman” nor is it a “female weakness.” Depressive illnesses are serious medical illnesses that affect more than 19 million American adults age 18 and over each year. Depression is a treatable medical illness that can occur in any woman, at any time, and for various reasons regardless of age, race or income.

### Prevalence

Approximately 12 million women in the United States experience clinical depression each year. About one in every eight women can expect to develop clinical depression during their lifetime. Depression occurs most frequently in women aged 25 to 44.

### Contributing Factors

Many factors in women may contribute to depression, such as developmental, reproductive, hormonal, genetic and other biological differences (e.g. premenstrual syndrome, childbirth, infertility and menopause). Social factors may also lead to higher rates of clinical depression among women, including stress from work, family responsibilities, the roles and expectations of women and increased rates of sexual abuse and poverty.

### Gender Differences

Women experience depression at roughly twice the rate of men. Girls 14-18 years of age have consistently higher rates of depression than boys in this age group. PMS/PMDD- Twenty to forty percent of women may experience premenstrual syndrome and an estimated 3 to 5 percent have symptoms severe enough to be classified as Premenstrual Dysphoric Disorder (PMDD).

### Marriage/Childbirth

Married people have a lower rate of depression than those living alone. However, unhappily married people have the highest rates of depression; happily married men have the lowest rates.

Approximately 10%-15% of all new mothers get postpartum depression, which most frequently occurs within the first year after the birth of a child.

### Co-occurring Illnesses

Research shows a strong relationship between eating disorders (anorexia and bulimia nervosa) and depression in women. About 90-95% of cases of anorexia occur in young females. Reported rates of bulimia nervosa vary from one to three out of 100 people. Research shows that one out of three depressed people also suffers from some form of substance abuse or dependence.

### Suicide

Men are more likely than women to die by suicide; women report attempting suicide approximately twice as often as men. An estimated 15 percent of people hospitalized for depression eventually take their own lives.

# What You Need to Know...



## **Treatment**

Depression in women is misdiagnosed approximately 30 to 50 percent of the time. Fewer than half of the women who experience clinical depression will ever seek care. Fortunately, clinical depression is a very treatable illness. More than 80 percent of people with depression can be treated successfully with medication, psychotherapy or a combination of both.

For more information, contact Mental Health America of Wisconsin at 414.276.3122, or via Email: [info@mhawisconsin.org](mailto:info@mhawisconsin.org).

Mental Health America is a United Way of Greater Milwaukee Agency.



[www.mhawisconsin.org](http://www.mhawisconsin.org)

---

600 West Virginia Street, Suite 502, Milwaukee, WI 53204 • P: 414.276.3122 • F: 414.276.3124  
133 S. Butler St., Room 330, Madison, WI 53703 • P: 608.250.4368 • F: 608.442.7907