

Depression in Older Adults

Depression affects more than 19 million Americans every year, regardless of age, race, or gender. While depression is not a normal part of the aging process, there is a strong likelihood of it occurring when other physical health conditions are present. For example, nearly a quarter of the 600,000 people who experience a stroke in a given year will experience clinical depression¹. Unfortunately, symptoms of depression are often overlooked and untreated when they coincide with other medical illnesses or life events that commonly occur as people age (e.g., loss of loved ones). However, clinical depression is never a “normal” response; it is a serious medical illness that should be treated at any age.

Prevalence

More than two million of the 34 million Americans age 65 and older suffer from some form of depression².

Co-occurring Illnesses

Symptoms of clinical depression can be triggered by other chronic illnesses common in later life, such as Alzheimer’s disease, Parkinson’s disease, heart disease, cancer and arthritis³.

Widowhood

One-third of widows/widowers meet criteria for depression in the first month after the death of their spouse, and half of these individuals remain clinically depressed after one year⁴.

Healthcare Costs

Older patients with symptoms of depression have roughly 50% higher healthcare costs than non-depressed seniors⁵.

Suicide

Depression is a significant predictor of suicide in elderly Americans². Comprising only 13% of the U.S. population, individuals aged 65 and older account for 20% of all suicide deaths, with white males being particularly vulnerable². Suicide among white males aged 85 and older (65.3 deaths per 100,000 persons) is nearly six times the suicide rate (10.8 per 100,000) in the U.S.².

Treatment

More than 55% of older persons treated for mental health services received care from primary care physicians. Less than 3% aged 65 and older received treatment from mental health professionals⁶. Primary care physicians accurately recognize less than one half of patients with depression, resulting in potentially decreased function and increased length of hospitalization⁷. Fortunately, clinical depression is a very treatable illness. More than 80% of all people with depression can be successfully treated with medication, psychotherapy or a combination of both⁸.

Older Adult Attitudes Toward Depression

According to a Mental Health America survey⁹ on attitudes and beliefs about clinical depression:

- Approximately 68% of adults aged 65 and over know little or almost nothing about depression.
- Only 38% of adults aged 65 and over believe that depression is a “health” problem.

What You Need to Know...



- If suffering from depression, older adults are more likely than any other group to “handle it themselves.” Only 42% would seek help from a health professional.
- Signs of depression are mentioned more frequently by people under age 64 than people aged 65 and over. These include “a change in eating habits” (29% vs. 15%), “a change in sleeping habits” (33% vs. 16%) and “sadness” (28% vs. 15%).
- About 58% of people aged 65 and older believe that it is “normal” for people to get depressed as they grow older.

References:

- 1) National Institute of Mental Health: “Co-occurrence of Depression With Stroke Fact Sheet.” Accessed Sept. 1999. Netscape: http://www.nimh.nih.gov/depression/co_occr/stroke.htm.
- 2) National Institute of Mental Health: “Older Adults: Depression and Suicide Fact Sheet.” Accessed August 1999. Netscape: <http://www.nimh.nih.gov/publicat/elderlydepsuicide.cfm>.
- 3) National Institute of Mental Health: “Depression Shares Symptoms With Other Medical Conditions.” Accessed September 1999. Netscape: <http://www.nimh.nih.gov/depression/senior/shares.htm>.
- 4) National Institute of Mental Health. “The Many Dimensions of Depression in Women: Women at Risk,” Accessed June 1999.
- 5) Netscape: <http://www.nimh.gov/depression/women/risk.htm>.
- 6) Unutzer, J., “Depressive symptoms and the cost of health services in HMO patients aged 65 years and older,” JAMA 277;20 (1997).
- 7) Olfson, M., Pincus, H.A., “Outpatient mental health care in nonhospital settings: distribution of patients across provider groups,”
- 8) American Journal of Psychiatry 153 (1996):1353-1356.
- 9) Sadosky, R., “Prevalence and recognition of depression in elderly patients,” American Academy of Family Physicians, 57;5 (1998):1096.
- 10) National Institute of Mental Health: “Depression: Treat it. Defeat it.” Accessed June 1999. Netscape: http://www.nimh.nih.gov/depression/genpop/gen_fact.htm.
- 11) National Mental Health Association, “American Attitudes about Clinical Depression and its Treatment,” (March 27, 1996).

Source: Mental Health America. Last updated: 11/08/2006.

www.mhawisconsin.org