

## Generalized Anxiety Disorder

Generalized Anxiety Disorder (GAD) is characterized by six months or more of chronic, exaggerated worry and tension that is unfounded or much more severe than the normal anxiety most people experience. People with this disorder usually expect the worst. They worry excessively about money, health, family or work, even when there are no signs of trouble. They are unable to relax and often suffer from insomnia. Many people with GAD also have physical symptoms, such as fatigue, trembling, muscle tension, headaches, irritability or hot flashes. Fortunately, through research supported by the National Institute of Mental Health (NIMH), effective treatments have been developed to help people with GAD.

### Prevalence

About 2.8 percent of the U.S. population ages 18 to 54 (about four million Americans) has GAD each year. GAD most often strikes people in childhood or adolescence, but it can begin in adulthood, too. It affects women more often than men.

### Causes

Some research suggests that GAD may run in families, and that it may grow worse during stress. GAD usually begins at an earlier age and symptoms may manifest themselves more slowly than in most other anxiety disorders.

### Treatments

**Medication:** Successful treatment may include antianxiety medications, such as buspirone (trade name BuSpar) and the benzodiazepines (Valium, Xanax, etc.) or antidepressants (Prozac, Paxil, etc.).  
**Behavioral therapy:** Behavioral therapy focuses on using specific relaxation techniques to change anxiety-causing behaviors. For example, one technique trains patients in a special breathing exercise involving slow, deep breaths to reduce anxiety. This is necessary because people who are anxious often hyperventilate, taking rapid, shallow breaths that can trigger rapid heartbeat, lightheadedness, and other symptoms. Another technique—exposure therapy—gradually exposes patients to what frightens them and helps them cope with their fears.

**Cognitive-behavioral therapy:** Like behavioral therapy, cognitive-behavioral therapy teaches patients to react differently to the situations and bodily sensations that trigger anxiety symptoms. However, patients also learn to understand how their thinking patterns contribute to their symptoms and how to change their thoughts so that symptoms are less likely to occur. This awareness of thinking patterns is combined with behavioral techniques to help people confront their feared situations.

### Co-occurring Illnesses

Research shows that GAD often coexists with depression, substance abuse, or other anxiety disorders. Other conditions associated with stress, such as irritable bowel syndrome, often accompany GAD. Patients with physical symptoms, such as insomnia or headaches, should also tell their doctors about their feelings of worry and tension. This will help the patient's health care provider to recognize that the person is suffering from GAD.

# What You Need to Know...



## Other Resources

Anxiety Disorders Association of America  
Phone Number: (240) 485 -1001  
Fax Number: (240) 485 -1035  
<http://www.adaa.org>

Freedom From Fear  
308 Seaview Avenue  
Staten Island, NY 10305  
Phone Number: (718) 351-1717  
Fax Number: (718) 980-5022  
[help@freedomfromfear.org](mailto:help@freedomfromfear.org)  
[www.freedomfromfear.com](http://www.freedomfromfear.com)

This fact sheet is a publication of Mental Health America's Campaign for America's Mental Health. The content of this fact sheet was adapted from material published by the National Institute of Mental Health.

Page last updated: 11/08/2006

[www.mhawisconsin.org](http://www.mhawisconsin.org)